

# Absence of Landau's Diamagnetism in Two Dimensions

S. FUJITA<sup>1</sup>(\*) and H. C. HO<sup>2</sup>(\*\*)

<sup>1</sup> *Department of Physics, University at Buffalo, SUNY  
Buffalo, New York 14260, USA*

<sup>2</sup> *Physics Division, National Center for Theoretical Sciences  
Hsinchu 30013, Taiwan*

PACS. 71.70.Di – Landau levels.

PACS. 73.20.-r – Electron states at surfaces and interfaces.

PACS. 71.10.Ca – Electron gas.

**Abstract.** – The statistical weight  $W$  as a function of energy  $E$  for quasielectrons with mass  $m^*$  subject to a fixed magnetic field  $\mathbf{B}$  is  $W/A = (m^*/\pi\hbar^2) E + (eB/\pi^2\hbar) \sum_{\nu=1}^{\infty} (-1)^\nu \nu^{-1} \cdot \sin(2\pi\nu E/\hbar\omega_c)$ , where  $A$  is the sample area, and  $\omega_c \equiv eB/m^*$  the cyclotron frequency. Significantly, there is no Landau's term proportional to  $B^2$  in 2D. This leads to the conclusion that the 2D electron system is always paramagnetic, but shows a magnetic oscillation.

Landau [1], Sondheimer and Wilson [2,3] discussed the de Haas-van Alphen (dHvA) oscillation [4] of a three-dimensional (3D) system of quasifree electrons. Nakamura [5] calculated the statistical weight  $W$  associated with the Landau states, and treated the dHvA oscillation. We extend Nakamura's theory to a 2D system.

Let us consider a dilute system of electrons, each with effective mass  $m^*$ , moving in a plane. Applying a magnetic field  $\mathbf{B}$  perpendicular to the plane, each electron will be in a Landau state of energy

$$E = (N_L + 1/2) \hbar\omega_c, \quad N_L = 0, 1, \dots, \quad (1)$$

where  $\omega_c = eB/m^*$  is the cyclotron frequency. The degeneracy of the *Landau level* (LL) is

$$\frac{eBA}{2\pi\hbar}, \quad A = \text{sample area.} \quad (2)$$

We introduce kinetic momenta

$$\Pi_x = p_x + eA_x, \quad \Pi_y = p_y + eA_y,$$

in terms of which the Hamiltonian  $\mathcal{H}$  for the electron is

$$\mathcal{H} = \frac{1}{2m^*} (\Pi_x^2 + \Pi_y^2) \equiv \frac{1}{2m^*} \Pi^2. \quad (3)$$

---

(\*) E-mail: fujita@buffalo.edu

(\*\*) E-mail: hcho@phys.cts.nthu.edu.tw




Fig. 1 – Quantization scheme for free electrons in 2D: without a magnetic field (dots), and in a magnetic field (circles).

After simple calculations, we obtain

$$dx d\Pi_x dy d\Pi_y = dx dp_x dy dp_y.$$

We can then represent quantum states by quasi phase-space elements  $dx d\Pi_x dy d\Pi_y$ . The Hamiltonian  $\mathcal{H}$  in eq. (3) does not depend on the position  $(x, y)$ . Assuming large normalization lengths  $(L_1, L_2)$ , we can represent the Landau states by concentric shells of the phase space having statistical weight  $2\pi \Pi \Delta \Pi \cdot L_1 L_2 (2\pi \hbar)^{-2} = eBA/2\pi \hbar$ , with  $A = L_1 L_2$  and  $\hbar \omega_c = \Delta (\Pi^2/2m^*) = \Pi \Delta \Pi/m^*$ . Hence, the LL degeneracy is given by eq. (2). Figure 1 represents a typical Landau state in the  $\Pi_x$ - $\Pi_y$  space.

As the field  $B$  is raised the separation  $\hbar \omega_c$  increases, and the quantum states are collected (or bunched) together. As a result of bunching, the density of states  $\mathcal{N}(\varepsilon)$  should change periodically since the Landau levels are spaced equally.

The electrons obey the Fermi-Dirac statistics. Considering a system of quasifree electrons, we define the Helmholtz free energy  $\mathcal{F}$  by

$$\mathcal{F} = N\mu - 2k_B T \cdot \sum_i \ln \left[ 1 + e^{(\mu - E_i)/k_B T} \right], \quad (4)$$

where  $\mu$  is the chemical potential, and the factor 2 arises from spin degeneracy. The chemical potential  $\mu$  is determined from the condition

$$\frac{\partial \mathcal{F}}{\partial \mu} = 0. \quad (5)$$

The magnetization  $\mathcal{M}$  for the system can be found from

$$\mathcal{M} = -\frac{1}{A} \frac{\partial \mathcal{F}}{\partial B}. \quad (6)$$

Equation (5) is equivalent to the usual condition that the number of electrons  $N$  can be obtained in terms of the Fermi distribution function  $F$ :

$$N = 2 \sum_i F(E_i), \quad F(E) \equiv \left[ e^{\beta(E-\mu)} + 1 \right]^{-1}. \quad (7)$$

The LL  $E_i$  is characterized by the Landau oscillator quantum number  $(N_L)_i$ .

Let us introduce the density of states  $dW/dE \equiv \mathcal{N}(E)$  such that  $\mathcal{N}(E)dE$  = number of states having energy between  $E$  and  $E + dE$ . We write eq. (4) in the form

$$\begin{aligned} \mathcal{F} &= N\mu - 2k_B T \cdot \int_0^\infty dE \frac{dW}{dE} \ln \left[ 1 + e^{(\mu-E)/k_B T} \right] \\ &= N\mu - 2 \int_0^\infty dE W(E) F(E). \end{aligned} \quad (8)$$

The statistical weight  $W$  is the total number of states having energies less than the Landau energy  $(N_L + 1/2)\hbar\omega_c$  in eq. (1). This  $W$  from fig. 1 is

$$W = \frac{L_1 L_2}{(2\pi\hbar)^2} 2\pi \Pi \Delta \Pi \cdot 2 \sum_{N_L=0}^\infty \Theta[E - (N_L + 1/2)\hbar\omega_c], \quad (9)$$

where  $\Theta(x)$  is the Heaviside step function

$$\Theta(x) = \begin{cases} 0 & \text{if } x < 0 \\ 1 & \text{if } x > 0 \end{cases}.$$

We introduce a dimensionless variable  $\varepsilon \equiv 2\pi E/\hbar\omega_c$ , and rewrite eq. (9) as

$$W(E) = C\hbar\omega_c \cdot 2 \sum_{N_L=0}^\infty \Theta[\varepsilon - (2N_L + 1)\pi], \quad (10)$$

with  $C = 2\pi m^* A (2\pi\hbar)^{-2}$ . We assume a high Fermi-degeneracy such that  $\mu \simeq \varepsilon_F \gg \hbar\omega_c$ . The sum in eq. (10) can be computed using Poisson's summation formula [6]

$$\sum_{n=-\infty}^\infty f(2\pi n) = \frac{1}{2\pi} \sum_{m=-\infty}^\infty \int_{-\infty}^\infty d\tau f(\tau) e^{-im\tau}, \quad (11)$$

where  $\sum_{n=-\infty}^\infty f(2\pi n + t)$ ,  $0 \leq t < 2\pi$  is by assumption uniformly convergent. We write the sum in eq. (10) as

$$2 \sum_{n=0}^\infty \Theta[\varepsilon - (2n + 1)\pi] = \Theta(\varepsilon - \pi) + \phi(\varepsilon; 0), \quad (12)$$

$$\phi(\varepsilon; x) \equiv \sum_{n=-\infty}^\infty \Theta(\varepsilon - \pi - 2\pi|n + x|).$$

Note that  $\phi(\varepsilon; x)$  is periodic in  $x$ . After the Fourier expansion, we set  $x = 0$  to obtain eq. (12). By taking the real part (Re) of eq. (12) and using eq. (11), we obtain

$$\text{Re} \{ \text{Eq. (12)} \} = \frac{1}{\pi} \int_0^\infty d\tau \Theta(\varepsilon - \tau) + \frac{2}{\pi} \sum_{\nu=1}^\infty (-1)^\nu \int_0^\infty d\tau \Theta(\varepsilon - \tau) \cos \nu\tau, \quad (13)$$

where we assumed  $\varepsilon \equiv 2\pi E/\hbar\omega_c \gg 1$ , and neglected  $\pi$  against  $\varepsilon$ . The integral in the first term in eq. (13) yields  $\varepsilon$ . The integral in the second term can be evaluated by integration by parts, and using  $d\Theta/dy = \delta(y)$ . We obtain

$$\int_0^\infty d\tau \Theta(\varepsilon - \tau) \cos \nu\tau = \frac{1}{\nu} \sin \nu\varepsilon.$$

Hence,

$$\text{Re}\{ \text{Eq. (12)} \} = \frac{1}{\pi}\varepsilon + \frac{2}{\pi} \sum_{\nu=1}^{\infty} \frac{(-1)^\nu}{\nu} \sin \nu\varepsilon. \quad (14)$$

Using eqs. (10) and (14), we obtain

$$W(E) = W_0 + W_{\text{osc}}, \quad (15)$$

$$W_0 = C\hbar\omega_c \frac{\varepsilon}{\pi} = A \frac{m^*E}{\pi\hbar^2} \quad (16)$$

$$W_{\text{osc}} = C\hbar\omega_c \frac{2}{\pi} \sum_{\nu=1}^{\infty} \frac{(-1)^\nu}{\nu} \sin \left( \frac{2\pi\nu E}{\hbar\omega_c} \right). \quad (17)$$

The oscillatory term  $W_{\text{osc}}$  contains an infinite sum with respect to  $\nu$ , but only the first term,  $\nu = 1$ , is important in practice as we see later. This term  $W_{\text{osc}}$  can generate magnetic oscillations. There is no term proportional to  $B^2$  generating the Landau diamagnetism. This is unexpected. We briefly discuss the difference between 2D and 3D systems.

In 3D, the LL  $E$  is given by

$$E = \left( N_L + \frac{1}{2} \right) \hbar\omega_c + \frac{p_z^2}{2m^*}.$$

The energy  $E$  is continuous in the bulk limit. The statistical weight  $W'$  is the total number of states having energies less than  $E$ . The allowed values of  $p_z$  are distributed over the range in which  $|p_z|$  does not exceed  $\{2m^*[E - (N_L + 1/2)\hbar\omega_c]\}^{1/2}$ . We obtain

$$W'(E) = C' \frac{(\hbar\omega_c)^{3/2}}{\sqrt{2\pi}} 2 \sum_{N_L=0}^{\infty} \sqrt{\varepsilon - (2N_L + 1)\pi}, \quad (18)$$

where  $C' \equiv V(2\pi m^*)^{3/2}/(2\pi\hbar)^3$ ,  $\varepsilon \equiv 2\pi E/\hbar\omega_c$ , and  $V \equiv L_1 L_2 L_3$  is the sample volume. We proceed similarly, and obtain

$$W'(E) = W'_0 + W'_L + W'_{\text{osc}}, \quad (19)$$

$$W'_0 = C' \frac{4}{3\sqrt{\pi}} E^{3/2} \quad (20)$$

$$W'_L = -C' \frac{1}{24\sqrt{\pi}} \frac{(\hbar\omega_c)^2}{E^{1/2}} \quad (21)$$

$$W'_{\text{osc}} = C' \frac{1}{\sqrt{2}} \left( \frac{\hbar\omega_c}{\pi} \right)^{3/2} \sum_{\nu=1}^{\infty} \frac{(-1)^\nu}{\nu^{3/2}} \sin \left( \frac{2\pi\nu E}{\hbar\omega_c} - \frac{\pi}{4} \right). \quad (22)$$

In detail, we write the sum in eq. (18) as

$$2 \sum_{n=0}^{\infty} \sqrt{\varepsilon - (2n+1)\pi} = (\varepsilon - \pi)^{1/2} + \psi(\varepsilon; 0), \quad (23)$$

$$\psi(\varepsilon; x) \equiv \sum_{n=-\infty}^{\infty} (\varepsilon - \pi - 2\pi|n+x|)^{1/2}.$$

Since  $\psi(\varepsilon; x)$  is periodic in  $x$ , we can use it for the Fourier expansion of eq. (23), and then set  $x = 0$ . By taking the real part (Re) of eq. (23) and using eq. (11), we obtain

$$\text{Re} \{ \text{Eq. (23)} \} = \frac{1}{\pi} \int_0^\varepsilon d\tau (\varepsilon - \tau)^{1/2} + \frac{2}{\pi} \sum_{\nu=1}^{\infty} (-1)^\nu \int_0^\varepsilon d\tau (\varepsilon - \tau)^{1/2} \cos \nu\tau, \quad (24)$$

where we neglected  $\pi$  against  $\varepsilon$ . The integral in the first term in eq. (24) yields  $(2/3)\varepsilon^{3/2}$ , leading to  $W'_0$  in eq. (20). The integral in the second term can be written after integrating by part, and changing the variable ( $\nu\varepsilon - \nu\tau = t$ ) as

$$\frac{1}{2\nu^{3/2}} \left[ \sin \nu\varepsilon \int_0^{\nu\varepsilon} dt \frac{\cos t}{\sqrt{t}} - \cos \nu\varepsilon \int_0^{\nu\varepsilon} dt \frac{\sin t}{\sqrt{t}} \right].$$

We now use asymptotic expansions for  $\nu\varepsilon = x \gg 1$ :

$$\int_0^x dt \frac{\sin t}{\sqrt{t}} \sim \sqrt{\frac{\pi}{2}} - \frac{\cos x}{\sqrt{x}} - \dots$$

$$\int_0^x dt \frac{\cos t}{\sqrt{t}} \sim \sqrt{\frac{\pi}{2}} + \frac{\sin x}{\sqrt{x}} - \dots$$

The second terms in the expansion lead to  $W'_L$  in eq. (21), where we used the identity

$$\sum_{\nu=1}^{\infty} \frac{(-1)^{\nu-1}}{\nu^2} = \frac{\pi^2}{12}.$$

The first terms lead to the oscillatory term  $W'_{\text{osc}}$  in eq. (22).

The term  $W'_0$ , which is independent of  $B$ , gives the weight equal to that for a free-electron system with no field. The term  $W'_L$  proportional to  $B^2$  is negative (diamagnetic), and can generate a Landau diamagnetic moment.

The energy  $E$  of the 3D system is continuous, and hence the system is manageable or soft. In contrast, the energy  $E$  of the 2D system is discrete, and hence the system is less manageable. This explains the absence of Landau diamagnetism for the 2D system.

The statistical weight  $W_{\text{osc}}$  in eq. (17) has a sine term. Hence, the density of states,  $\mathcal{N} = dW/dE$ , has an oscillatory part of the form

$$\sin \left( \frac{2\pi E}{\hbar\omega_c} \right), \quad E \equiv \frac{\Pi^2}{2m^*}.$$

If the density of states oscillates violently in the drop of the Fermi distribution function:  $F(E) \equiv [e^{\beta(E-\mu)} + 1]^{-1}$ , the delta-function replacement formula

$$-\frac{dF}{dE} = \delta(E - \mu),$$

cannot be used. The width of  $-dF/dE$  is of the order  $k_B T$ . The critical temperature  $T_c$  below which oscillations can be observed is  $k_B T_c \sim \hbar \omega_c$ . For  $T < T_c$ , we may proceed as follows. Let us consider the integral

$$I = \int_0^\infty dE F(E) \sin(2\pi E/\hbar \omega_c).$$

We introduce a new variable  $\zeta \equiv \beta(E - \mu)$ , and extend the lower limit to  $-\infty$  (low-temperature limit) so that

$$\int_0^\infty dE \cdots \frac{1}{e^{\beta(E-\mu)} + 1} = \frac{1}{\beta} \int_{-\mu\beta}^\infty d\zeta \cdots \frac{1}{e^\zeta + 1} \rightarrow \frac{1}{\beta} \int_{-\infty}^\infty d\zeta \cdots \frac{1}{e^\zeta + 1}.$$

With the help of the standard integral

$$\int_{-\infty}^\infty d\zeta \frac{e^{ia\zeta}}{e^\zeta + 1} = \frac{\pi}{i \sinh \pi a},$$

we obtain

$$I = -\frac{\pi k_B T \cos(2\pi \varepsilon_F/\hbar \omega_c)}{\sinh(2\pi^2 k_B T m^*/\hbar e B)}. \quad (25)$$

For very low fields the oscillation number in the range  $k_B T$  becomes great, and hence the sinusoidal contribution must cancel out. This effect is represented by the factor

$$[\sinh(2\pi^2 k_B T m^*/\hbar e B)]^{-1}.$$

We calculate the free energy indicated in eq. (8) using the statistical weight  $W$  in eq. (15), and obtain

$$\mathcal{F} = N\mu - 2A \frac{m^*}{\pi \hbar^2} \varepsilon_F + 2A \frac{eB}{\pi \hbar} k_B T \cdot \sum_{\nu=1}^{\infty} \frac{(-1)^\nu}{\nu} \frac{\cos(2\pi \nu \varepsilon_F/\hbar \omega_c)}{\sinh(2\pi^2 \nu k_B T m^*/\hbar e B)}, \quad (26)$$

where we used the integration formula (25), and took the low-temperature limit except for the oscillatory terms. The magnetization  $\mathcal{M}$  can be obtained using eq. (6).

So far, we have not considered the Pauli spin magnetization [7]

$$\mathcal{M}_{\text{Pauli}} = 2\mu_B^2 B \mathcal{N}_0(\varepsilon_F)/A = 2n\mu_B^2 B/\varepsilon_F,$$

with  $\mu_B$  (Bohr magneton)  $= e\hbar/2m$  and  $n$  (electron number density)  $= m^* \varepsilon_F/\pi \hbar^2$ . Using eqs. (6), (7) and (26), we obtain the total magnetization  $\mathcal{M}_{\text{tot}} = \mathcal{M}_{\text{Pauli}} + \mathcal{M}_{\text{osc}}$ :

$$\mathcal{M}_{\text{tot}} = 2n\mu_B \frac{\mu_B B}{\varepsilon_F} \left[ 1 + 2 \frac{k_B T}{\mu_B B} \frac{m}{m^*} \frac{\cos(2\pi \varepsilon_F/\hbar \omega_c)}{\sinh(2\pi^2 k_B T m^*/\hbar e B)} \right]. \quad (27)$$

The  $B$ -dependence of  $\mathcal{F}$  is contained in the last term in eq. (26). The linear  $B$ -dependence of the multiplication factor is much stronger than the  $B$ -dependence of the alternating series. Therefore, the contribution from the  $B$ -derivative of the series is neglected. The variation of the statistical weight  $W$  is periodic in  $B^{-1}$ , but it is far from sinusoidal. Only the first oscillatory term,  $\nu = 1$ , is important and kept in eq. (27) since  $\sinh(2\pi^2 k_B T m^*/\hbar e B) \gg 1$ . The width of  $dF/dE$  is finite for a finite  $T$ . In this  $E$ -range, many oscillations can occur if the field  $B$  is made low. We assumed this condition to obtain eq. (27). The magnetic susceptibility  $\chi$  is defined by the ratio  $\chi = \mathcal{M}/B$ .

In conclusion, the 2D system is intrinsically paramagnetic since the Landau's diamagnetic term is absent, but the system exhibits a dHvA oscillation.

## REFERENCES

- [1] LANDAU L. D., *Z. Physik*, **64** (1930) 629.
- [2] SONDHEIMER E. H. AND A. H. WILSON, *Proc. Roy. Soc. London, Ser. A*, **210** (1951) 173.
- [3] WILSON A. H., *The Theory of Metals* (Cambridge University Press, London) 1954, p. 676.
- [4] DE HAAS W. J. AND P. M. VAN ALPHEN, *Leiden Comm.*, (1930) 208d; (1930) 212a; (1932) 220d.
- [5] NAKAMURA T., *Magnetism* (Makishoten, Tokyo) 1965, p. 75–82.
- [6] COURANT R. AND D. HIBERT, *Methods of Mathematical Physics*, Vol. **1** (Interscience-Wiley, New York) 1953, p. 76–77.
- [7] PAULI W., *Z. Physik*, **41** (1927) 81.